

2019-08-13

Dietitian-led clinics in primary care: A scoping review

Hickson, Mary

<http://hdl.handle.net/10026.1/14839>

10.11124/jbisrir-d-19-00025

The JBI Database of Systematic Reviews and Implementation Reports

Lippincott, Williams & Wilkins

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Dietitian-led clinics in primary care: A scoping review protocol

Authors

Mary Hickson^{1,2,3}

Amanda Wanner^{2,3,4}

Avril Collinson^{1,3}

¹Institute of Health and Community, University of Plymouth, Plymouth, UK

²NIHR Collaboration for Leadership in Applied Health Research and Care South West Peninsula (PenCLAHRC), University of Plymouth, Plymouth, UK

³University of Plymouth Centre for Innovations in Health and Social Care: A Joanna Briggs Institute Centre of Excellence, Plymouth, UK

⁴Faculty of Medicine and Dentistry, University of Plymouth, Plymouth, UK

Corresponding Author:

Mary Hickson

mary.hickson@plymouth.ac.uk

Funding

This review is funded by a project grant award to the University of Plymouth by the British Dietetic Association (BDA) General Education Trust (a Charitable body specifically convened to support dietetic research and development projects). Officers of the BDA and trustees of the General Education Trust had no role in the development of the protocol, design of the study, data collection and collation, nor interpretation and discussion of the data. Officers of the BDA did assist with identification of grey literature through their own knowledge and use of professional networks.

This research was also supported in part by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care South West Peninsula. The views

expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Conflicts of Interest

Mary Hickson and Avril Collinson are both members of the British Dietetic Association (funder) and registered dietitians. Amanda Wanner declares no conflict of interest.

Abstract

Review objective and questions: The objective of this scoping review is to examine and map the existing evidence exploring or evaluating the implementation, cost and/or effectiveness of dietitian-led clinics in primary care.

- Does a dietitian-led clinic in general practice and/or primary care improve patient satisfaction and clinical outcomes?
- Does a dietitian-led clinic in general practice and/or primary care reduce costs?
- Is it feasible to implement dietitian-led clinics in general practice and/or primary care?

Introduction

The World Health Organization (WHO) describes primary care as “first-contact, accessible, continued, comprehensive and coordinated care. First-contact care is accessible at the time of need; ongoing care focuses on the long-term health of a person rather than the short duration of the disease; comprehensive care is a range of services appropriate to the common problems in the respective population and coordination is the role by which primary care acts to coordinate other specialists that the patient may need”.^{1(para.3)} The terms ‘general practice’ and ‘family medicine’ are both synonymous with primary care and may be used interchangeably in the literature. How primary care is organized varies between countries and may be centred on the general practitioner (GP) (or primary care physician) or take a more team orientated approach recognising the primary care team (or family medicine team). The WHO describes the primary care team as “a group of fellow professionals with complementary contributions to make in patient care”,^{1(para.5)} of which a dietitian may be one member.

Dietitians have a skill-set that enables them to lead on the therapeutic support provided to patients with certain conditions that are amenable to treatment with dietary manipulation. Examples of such conditions include diabetes mellitus, cardio-vascular disease, over- and under-weight, food allergies, chronic obstructive pulmonary disease, gastrointestinal, renal and liver conditions. Dietitians have historically worked largely in acute hospital settings² however, there is little information on dietitians who work in primary care. This may be a service commissioned by the general practice to private dietitians, or contracted from dietetic services based in the acute or community sectors.

Throughout the developed world healthcare is changing. Some of the contributing factors include demographic shifts such as the ageing population, the increase of long-term conditions, increase of dementia, changes in the diversity of society, health inequalities and limited funding.² This has led to

an increased demand within the primary healthcare sector at the same time as GP numbers in many countries are declining, including the UK,³ USA,⁴ and Australia.⁵

Given the skill set of the dietitian it may be that new models of care with the primary care setting, could see dietitians taking a lead in delivering primary care for relevant diagnosed patients and providing cheaper, more efficient and effective service in comparison to the traditional approach of GP support and referral for specialist treatment. Indeed, this has been promoted by the UK government as a way to tackle work pressures within primary care and general practice.^{6, 7}

A dietitian-led clinic is any clinic run and managed by a registered dietitian, and in this scoping review is limited to the primary care setting, where the clinic is likely to support the work of GPs. This would mean that patients with relevant diagnoses (see above) could be referred by another healthcare professional, self-refer or be invited to the clinic for diet and lifestyle advice and support.

There are several systematic reviews that indicate how advice provided by a dietitian can improve outcomes in specific conditions, such as hypertension,⁸ diabetes, weight loss and diet quality.⁹ The evidence for gestational weight gain⁹ and prevention of gestation diabetes¹⁰ is weaker primarily due to lower quality study design. Other systematic reviews have explored interventions to manage weight in children¹¹ and adults,¹² type 2 diabetes,¹³ diabetes prevention,¹⁴ and Mediterranean diet and healthy eating,¹⁵ but these studies were not specific to dietetic interventions, although they included studies examining dietetic care. They all showed that dietary interventions could improve outcomes, and some showed that care provided by dietitians achieved superior outcomes, but the quality of the study designs were often weak. Other original studies also support the view that dietitians and/or dietary counselling (which dietitians are uniquely trained to deliver) are effective in improving clinical outcomes in a number of health conditions.¹⁶⁻²⁰ Therefore, it would seem that greater utilization of dietetic interventions in the primary care setting could be an effective way to manage many common chronic diseases, however, it is important to demonstrate that interventions are effective in the setting in which they will be delivered.

A review by Mitchell et al.⁹ is the only one available looking specifically at dietitians in primary care, and this included only randomised controlled trials. They did not search for any particular disease category but looked at any patient receiving dietetic consultations. The conditions treated included HIV, cardiovascular disease, obesity, hypertension, diabetes, impaired fasting glucose, gestational diabetes and colorectal cancer. The results show fair (Grade 2) evidence for dietetic consultations for adults in primary care settings for improvement in diet quality, diabetes outcomes (including blood glucose and glycated haemoglobin values), and weight loss outcomes (e.g. changes in weight and waist circumference) and to limit gestational weight gain. The evidence for controlling lipid levels and blood pressure is limited (Grade 3), but this review included only studies where the provision of nutritional care was exclusively by a dietitian. Many of the studies testing interventions for cardiovascular diseases have multi-disciplinary team interventions and these, with the dietetic contribution, would not have been included in this review.

Thus, there is some evidence for the efficacy of dietetic care in primary care. Nevertheless, there is a lack of information concerning the broader contribution dietitians may make within the primary care

setting, including cost effectiveness and the range of conditions that dietitians may successfully manage. There may also be useful qualitative information as well as quantitative work. A preliminary search of the *JBI Database of Systematic Reviews and Implementation Reports*, Cochrane Library, PubMed and CINAHL databases found no scoping reviews exploring dietetic care with the primary care setting. A search of the PROSPERO database found no similar systematic review protocol registered or ongoing.

Objective

The objective of this scoping review is to examine and map the existing evidence exploring or evaluating the implementation, cost and/or effectiveness of dietitian-led clinics in primary care.

Review questions

Does a dietitian-led clinic in general practice and/or primary care improve patient satisfaction and clinical outcomes?

Does a dietitian-led clinic in general practice and/or primary care reduce costs?

Is it feasible to implement dietitian-led clinics in general practice and/or primary care?

Keywords

Dietitian-led; primary care; general practice;

Inclusion criteria

Participants

The review will consider studies that include dietitian or nutritionist-led clinics treating patients with any conditions. Both terms will be considered because the professions are linked and the name varies between countries. However, to be comparable any dietitian or nutritionist-led studies would need to require the dietitian or nutritionist to have formal accreditation.

Concept

The proposed review is designed to explore the feasibility, organization and effectiveness of dietitian-led clinics within a primary care setting. Therefore, all studies with a focus on any aspect of dietitian or nutritionist led healthcare services for any disease group will be considered. Of particular interest will be any evidence of cost effectiveness in comparison to the usual organization of services.

Context

The context for this review will be primary care or general practice. General practice is part of primary care, but both terms will be of interest since services provided as part of primary care will be of interest even if not based in general practice. Both terms may be used interchangeably in papers and so it is important to identify all sources of evidence. Dietitian or nutritionist-led clinics in hospitals,

regional healthcare facilities or specialist centers will not be included. Where studies have been conducted in the community, they will be relevant if recruitment has included general practitioners.

This review will also only consider evidence from developed countries, since the settings are more likely to be comparable. It is recognised that healthcare is delivered and organized differently even in developed countries, but findings from developing countries will have less applicability. The World Bank country classifications will be used to decide which countries are deemed developing.²¹

Types of studies

This scoping review will consider all available publications that have a focus on dietitian or nutritionist led clinical care in a primary care setting. These may include experimental, quasi-experimental, observational and qualitative studies. Systematic reviews will be considered, as well as text and opinion papers, case studies, and relevant academic presentations, in both peer-reviewed and grey literature. Dietetic networks will be used to identify relevant grey literature from other countries.

Methods

The proposed systematic review will be conducted in accordance with the Joanna Briggs Institute methodology for scoping reviews.²²

Search strategy

The search strategy aims to find both published and unpublished studies. The systematic search will be developed and run by an experienced information specialist (AW). The initial strategy was iteratively designed by testing search terms against a pre-defined list of relevant articles and tested in several different databases. The final strategy will be translated for use in each of the databases (an example of the Ovid MEDLINE search is in Appendix 1). The searches will be limited to the last 10 years, excluding studies prior to 2008. Limiting the search to the last 10 years ensures that the information retrieved will be as relevant as possible to today's healthcare setting. There will be no limit on language applied to the searches. The following databases will be searched: MEDLINE (Ovid), Embase (Ovid), PsycINFO (Ovid), CINAHL (Ebsco), AMED (Ebsco), British Nursing Index (Proquest), and Cochrane Library (Wiley). Next unpublished studies will be sought through requests to experts and professional bodies using existing dietetic networks, and through searching Open Grey, ClinicalTrials.gov and EU Clinical Trials Register. Finally, the reference lists of each of the included papers will be hand searched to identify any further studies.

Study selection

Following the search, all identified citations will be collated and uploaded into EndNote X8.2 (Clarivate Analytics, PA, USA) and duplicates removed. The set will then be uploaded to Rayyan QCRI²³ and titles and abstracts screened by two independent reviewers for assessment against the inclusion criteria for the review. Any disagreements will be solved by consensus or by the decision of a third reviewer. The full text of studies that may meet the inclusion criteria will be retrieved and re-screened to confirm inclusion. Full text studies that do not meet the inclusion criteria will be excluded and

reasons for exclusion will be provided in an appendix in the final systematic review report. The final full text papers will be imported into JBI System for the Unified Management, Assessment and Review of Information (SUMARI).

Papers will be included if the setting is based in primary care and general practice; it is regarding health service delivery in a developed country; it is about dietetic or nutritionist led clinics, consultations, advice or counselling; and the paper was published in or after 2008. Studies testing the efficacy of a nutrient, food or dietary pattern but involves a dietitian to deliver information will be excluded.

Data extraction

Data will be extracted from the included papers by two independent reviewers using an adapted version of the Joanna Briggs Institute results extraction instrument.²⁴ The data extracted will include specific details about the population, concept, context, study methods and key findings relevant to the review objective. This information will be tabulated including the following; author/s, year of publication, country, setting, purpose of the study, study design, intervention (where relevant), participants, relevant outcomes such as cost efficacy or relevant clinical outcome data, and key findings that related to the review question. The draft results extraction instrument will be tested on the first five papers and modified as necessary, further revisions may be made during the process of extracting data from the remaining studies. Modifications will be detailed in the full scoping review report. Any disagreements that arise between the reviewers will be resolved through discussion, or with a third reviewer. Authors of papers will be contacted to request missing or additional data, where required.

Presentation of the results

The extracted data will be presented in diagrammatic or tabular form in a manner that aligns with the objective of this scoping review. A narrative summary will accompany the tabulated and/or charted results and will describe how the results relate to the review's objective and question/s.

References

1. World Health Organization. Primary Health Care: Main Terminology [internet]. Geneva: World Health Organization; 2019 [cited 2019 Jan 21] Available from: <http://www.euro.who.int/en/health-topics/Health-systems/primary-health-care/main-terminology>.
2. Hickson M, Child J, Collinson A. Future Dietitian 2025: informing the development of a workforce strategy for dietetics. *J Hum Nutr Diet*. 2018;31(1):23-32.
3. NHS Digital. General and Personal Medical Services, England As at 30 September 2017, Provisional Experimental statistics [internet]. NHS Digital; 2017 [cited 2019 Jan 21] Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/general-and-personal-medical-services/as-at-30-september-2017-provisional-experimental-statistics>.

- 208 4. Dall T, West T, Chakrabarti R, Reynolds R, Iacobucci W. The Complexities of Physician Supply
209 and Demand: Projections from 2016 to 2030. Washington, DC, USA: Association of American
210 Medical Colleges; 2018.
- 211 5. Government of Western Australia. General practice workforce supply and training in Western
212 Australia: Optimising Western Australia's prevocational training to support general practice
213 workforce development. Department of Health; 2018.
- 214 6. NHS England. General practice forward view. London (UK): NHS; 2016.
- 215 7. NHS. The NHS Long Term Plan – a summary. London (UK): NHS; 2019.
- 216 8. Riegel GR, Ribeiro PAB, Rodrigues MP, Zuchinali P, Moreira LB. Efficacy of nutritional
217 recommendations given by registered dietitians compared to other healthcare providers in
218 reducing arterial blood pressure: Systematic review and meta-analysis. *Clin Nutr.*
219 2018;37(2):522-531.
- 220 9. Mitchell LJ, Ball LE, Ross LJ, Barnes KA, Williams LT. Effectiveness of Dietetic Consultations in
221 Primary Health Care: A Systematic Review of Randomized Controlled Trials. *J Acad Nutr Diet.*
222 2017;117(12):1941-1962.
- 223 10. Oostdam N, van Poppel MN, Wouters MG, van Mechelen W. Interventions for preventing
224 gestational diabetes mellitus: a systematic review and meta-analysis. *J Womens Health*
225 (Larchmt). 2011;20(10):1551-1563.
- 226 11. Ho M, Jensen ME, Burrows T, Neve M, Garnett SP, Baur L, et al. Best practice dietetic
227 management of overweight and obese children and adolescents: A 2010 update of a systematic
228 review. *JBPI Database of Systematic Reviews and Implementation Reports.* 2013;11(10):190-293.
- 229 12. Flodgren G, Deane K, Dickinson HO, Kirk S, Alberti H, Beyer FR, et al. Interventions to change the
230 behaviour of health professionals and the organisation of care to promote weight reduction in
231 overweight and obese people. *Cochrane Database Syst Rev.* 2010.
- 232 13. Moller G, Andersen HK, Snorgaard O. A systematic review and meta-analysis of nutrition
233 therapy compared with dietary advice in patients with type 2 diabetes. *Am J Clin Nutr.*
234 2017;106(6):1394-1400.
- 235 14. Sun Y, You W, Almeida F, Estabrooks P, Davy B. The Effectiveness and Cost of Lifestyle
236 Interventions Including Nutrition Education for Diabetes Prevention: A Systematic Review and
237 Meta-Analysis. *J Acad Nutr Diet.* 2017;117(3):404-421.e436.
- 238 15. Maderuelo-Fernandez JA, Recio-Rodriguez JI, Patino-Alonso MC, Perez-Arechaederra D,
239 Rodriguez-Sanchez E, Gomez-Marcos MA, et al. Effectiveness of interventions applicable to
240 primary health care settings to promote Mediterranean diet or healthy eating adherence in
241 adults: A systematic review. *Prev Med.* 2015;76:S39-55.
- 242 16. Canani RB, Leone L, D'Auria E, Riva E, Nocerino R, Ruotolo S, et al. The effects of dietary
243 counseling on children with food allergy: a prospective, multicenter intervention study. *J Acad*
244 *Nutr Diet.* 2014;114(9):1432-1439.
- 245 17. Coppel KJ, Kataoka M, Williams SM, Chisholm AW, Vorgers SM, Mann JI. Nutritional
246 intervention in patients with type 2 diabetes who are hyperglycaemic despite optimised drug
247 treatment--Lifestyle Over and Above Drugs in Diabetes (LOADD) study: randomised controlled
248 trial. *BMJ.* 2010;341:c3337.
- 249 18. Lin JS, O'Connor E, Whitlock EP, Beil TL, Zuber SP, Perdue LA, et al. Behavioral Counseling to
250 Promote Physical Activity and a Healthful Diet to Prevent Cardiovascular Disease in Adults:
251 Update of the Evidence for the U.S. Preventive Services Task Force. 2010.

19. Sialvera TE, Papadopoulou A, Efstathiou SP, Trautwein EA, Ras RT, Kollia N, et al. Structured advice provided by a dietitian increases adherence of consumers to diet and lifestyle changes and lowers blood low-density lipoprotein (LDL)-cholesterol: the Increasing Adherence of Consumers to Diet & Lifestyle Changes to Lower (LDL) Cholesterol (ACT) randomised controlled trial. *J Hum Nutr Diet.* 2018;31(2):197-208.
20. Sugawara N, Sagae T, Yasui-Furukori N, Yamazaki M, Shimoda K, Mori T, et al. Effects of nutritional education on weight change and metabolic abnormalities among patients with schizophrenia in Japan: A randomized controlled trial. *J Psychiatr Res.* 2018;97:77-83.
21. World Bank. Developing Countries [internet]. The Hague: International Statistical Institute; 2018 [cited 2019 Jan 21]. Available from: <https://www.isi-web.org/index.php/resources/developing-countries>.
22. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implementation Science.* 2010;5:69-69.
23. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev.* 2016;5(1):210.
24. Peters MDJ, Godfrey C, McInerney P, Baldini Soares C, Khalil H, Parker D. Chapter 11: Scoping Reviews. In: Aromataris E, Munn Z (Editors). *Joanna Briggs Institute Reviewer's Manual*. The Joanna Briggs Institute, 2017. Available from: <https://reviewersmanual.joannabriggs.org/>

Appendix 1

Example search for Ovid MEDLINE

Searches

- 1 dietetics/ or nutritionists/ or nutrition assessment/ or (dieti?ian* or dietetic*).ti,ab,kw.
- 2 (counse?ling or advice or consultation* or intervention).ti,ab,kw. or counseling/
- 3 ((diet* or nutrition*) adj (counsel?ing or advice or consultation*)).ti,ab,kw.
- 4 (1 and 2) or 3
- 5 limit 4 to yr="2008 -Current"